

# 2018

1.

(ZY/T001. 1-94) [1]

1

2

3

4

X

1

4

2.

2009

1999

2014

8

[2-4]

48

1

38

2

3

(ZY/T001. 1-94)

2017

2017

[1, 5, 6]

1.

2.

3.

4.

5.

6.

1.

2.

3.

4.

5.

6.

1.

0.5~1

2.

3.

4.

2h

1 7 1

5.

8

1999

2014

[3-4]

1.

/

( )

2.

3.

1.

2.

3.

4.

5.

6.

1.

(ZY/T001.1-94)

[1]

90%

60%

90%

30%

60%

30%

2.

4

0

1

2

3

=

-

/

× 100%

3.

clinical pulmonary infection score CPIS [7]

CPIS

X

1

CPIS

	0	1	2
(12 )	36 38	38 39	39 36
× 10 <sup>9</sup> /L	4 11	11 17	4 17
(24 )			
(PaO <sub>2</sub> /Fi O <sub>2</sub> , kPa)	33		33
X			

2

2

	0 37.5
	1 37.5~38
	2 38~39
	3 39
	0
	1
	2
	3
	0
	1 10~50ml
	2 50~100ml
	3 100ml
	0
	1
	2
	3
	0
	1

	0					
	1					
	2					
	3					
X	0					
	1					
	2					
	0					
	1	WBC	10	10 <sup>9</sup> /L	4	10 <sup>9</sup> /L
	2	WBC	10	10 <sup>9</sup> /L	4	10 <sup>9</sup> /L

- [1] . . . . . (ZY/T001.1-94) [M]. : , 1994: 3-4
- [2] . . . . . [M]. : , 2009: 38-44
- [3] . . . . . [J]. , 1999, 22(4): 201-203
- [4] , . . . . . 8 [M]. : , 2014: 42
- [5] . . . . . [M]. , 2017: 173-181
- [6] . . . . . [M]. , 2017: 142-146
- [7] LUNA CM, BLANZACO D, NIEDERMAN MS, et al. Resolution of ventilator-associated pneumonia: prospective evaluation of the clinical pulmonary infection score as an early clinical predictor of outcome [J]. CritCareMed, 2003, 31(3): 676-682.