

2018

NCCN

2017

[1]

1.

1

1

2.

1

B

CT MRI PET-CT

2

B CT

3

CA125 HE4 CEA CA199 AFP

2014

[2]

1.

2.

3.

4.

5.

6.

1.

() ()

2.

3.

()

4.

()

5.

6.

()

1.

1

90% 500ml 3

2

15g 9cm 12cm

5mm

1

2

3

9cm 12cm

5mm

1

2

4

8h

2.

1 30min 10 1
2/15Hz 20 30min

3.

NCCN

2017

1.

2.

3.

4.

5.

2002

WHO

1.

4

0 1 2 3

1

/
:
:
:
70%
30%
30%

= +

2.

KPS

: 10
:
: 10
: 10

3.

1

CR 4
PR 30% 4
PD 20%
SD PR PD

2

CR
/ IR/SD /
PD /

1.

1

2.

KPS 2

3.

RECIST

3

4.

[1]NCCN Clinical Practice Guidelines in Oncology- ovarian cancer(2017 Version I)[DB/OL].<http://www.nccn.org>

[2] . [M]. 2014 448-464.

1.

	1	2	3
	2kg	5kg	10kg
		1/3	1/2
		0.5 1	1
		1 2	3
	2 3 /	4 5 / 1 2 /	3 /

KPS

Karnofsky KPS	
100	
90	
80	
70	
60	
50	
40	
30	
20	
10	
0	

3

RECIST

1

a.

20mm CT

10mm

b.

<20mm CT

<10mm)

/

4

a. MRI CT

CT MRI 10mm

CT 5mm

b.

c.

CR

d.

CR

2

/

a. 5
10

b.

a.
 CR 4
 PR 30% 4
 SD stabile di sease PR
 PD
 PD 20%

b.
 CR
 IR/SD incomplete response /
 /
 PD /